

Hypoglossal Neurinoma of The Carotid Space Segment

Natalia Rotaru, MD, PhD¹, Maxim Crivcheanscii, MD¹, Janna Punga, MD, PhD^{1,2}, Eugeniu Condrea, MD³, and Ion Codreanu, MD, PhD^{1*}

¹Department of Radiology and Medical Imaging, State University of Medicine and Pharmacy "Nicolae Testemitanu", Chisinau, Republic of Moldova

²Institute of Oncology, Chisinau, Republic of Moldova

³Institute of Neurology and Neurosurgery, Chisinau, Republic of Moldova

Received October 16, 2016; Accepted October 21, 2016; Published October 27, 2016

Keywords: Hypoglossal nerve palsy, Neurinoma, MRI.

A 33-year-old female presented with deviation of her tongue towards the right side (**Figure 1, A**). Magnetic resonance imaging (MRI) of the head revealed a 1.0 x 0.8cm mass abutting the right internal carotid artery in close proximity to the hypoglossal canal. The mass proved relatively isointense to brain parenchyma on T1 weighted images (**Figure 1, B**), slightly hyperintense on T2 weighted images (**Figure 1, C**), with a higher signal intensity on T2-tirm imaging (**Figure 1, D**). A "T2 shine through" phenomenon was also noted on Diffusion Weighted Imaging (DWI) with Apparent Diffusion Coefficient (ADC) mapping (**Figure 1, E and F**), the findings being consistent with a hypoglossal neurinoma (schwannoma) of the carotid space segment.




Figure 1. Magnetic resonance imaging of the head. Coronal T2-weighted imaging (A) shows rightward-deviating tongue (asterisks). Axial T1-weighted (B), T2-weighted (C) and T2-tirm (D) imaging show a mass (pointed by arrows) distally to the hypoglossal canal (pointed by a curved arrow). Increased signal intensity on both DWI (E) and ADC imaging (F) is consistent with a "T2 shine through" phenomenon (not restricted diffusion).

The hypoglossal nerve is divided into five segments: the medullary, cisternal, skull base, carotid space, and sublingual segments. Because each segment is usually affected by different disorders, localizing a lesion to a particular segment allows the physician to narrow the differential diagnosis [1]. Cranial nerve schwannomas are usually isolated lesions, except when they are associated with neurofibromatosis type 2 (abnormality of chromosome 22) [2]. Because of the substantial chance of nerve palsy following resection, obtaining an accurate preoperative diagnosis with the identification of the involved structures is crucial to the management strategy [3]. MRI is the investigation of choice in the diagnosis of schwannoma and the identification of the nerve of origin [3]. Evolving MRI techniques such as super selective diffusion tensor tractography (DTT) can provide further details related to the tumor and cranial nerve interface, including the presence or absence of penetrating fibers [4].

Corresponding author: Ion Codreanu, M.D., Ph.D., Department of Radiology and Medical Imaging, State University of Medicine and Pharmacy "Nicolae Testemitanu", Chisinau, MD 2025 Republic of Moldova, Phone: (373) 6769 3616, E-mail: codrion@yahoo.com.

Citation: Rotaru N, Crivcheanschii M, Punga J, Condrea E & Codreanu I (2016) Hypoglossal Neurinoma of The Carotid Space Segment. *Int J Med Clin Imaging*, 1(1): 8-9.

Copyright: ©2016 Rotaru N, Crivcheanschii M, Punga J, Condrea E & Codreanu I. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

REFERENCES

1. Thompson EO, Smoker WR. Hypoglossal nerve palsy: a segmental approach. *Radiographics*. 1994;14(5):939-958.
2. Spilberg G, Marchiori E, Gasparetto EL, Cabral RF, Takayassu TC, Batista RR, Vieira IG. Magnetic resonance findings of neurofibromatosis type 2: a case report. *Cases J*. 2009;2:6720
3. Liu HL, Yu SY, Li GK, Wei WI. Extracranial head and neck Schwannomas: a study of the nerve of origin. *Eur Arch Otorhinolaryngol*. 2011;268(9):1343-1347.
4. Wei PH, Qi ZG, Chen G, Hu P, Li MC, Liang JT, Guo HC, Ling F, Bao YH. Identification of cranial nerves near large vestibular schwannomas using superselective diffusion tensor tractography: experience with 23 cases. *Acta Neurochir (Wien)*. 2015;157(7):1239-1249.