

## 281. ANTIBIOTIC SUSCEPTIBILITY OF ENTEROBACTERIACEAE STRAINS ISOLATED FROM URINARY TRACT INFECTIONS

**Carlos Tortosa Valiente, Lukas Eckardt**

Scientific adviser: Székely Edit, MD, PhD, University of Medicine and Pharmacy Targu Mures, Romania

Enterobacteriaceae are the most frequent causes of urinary tract infections. We analysed the antibiotic susceptibility of urinary isolates recovered at the Microbiology Laboratory of Mures County Emergency Clinical Hospital.

**Materials and methods:** We collected data from the electronic data base of the laboratory. All successive non-duplicate, clinically significant Enterobacteriaceae strains tested during a one year period (2015) were included in our study. Recurrent isolates were considered for analysis only if there were phenotypically different. Pluribacterial samples were excluded from the study.

**Results:** A total of 672 strains from 651 patients were involved in our study. The most frequent was *E. coli* (n=500, 74%), followed by *Klebsiella pneumoniae* (90, 13%), *Proteus mirabilis* (34, 5%), *Serratia marcescens* (18, 3%) and others (5%). The highest susceptibility was registered for ertapenem (93%). The least active antibiotic was ampicillin (31%). Relatively low susceptibility was detected against fluoroquinolones (64%) and trimethoprim-sulfamethoxazol (60%). In case of *E. coli* the highest susceptibility was registered for ertapenem (99.8%) and nitrofurantoin (99%). In case of *Klebsiella pneumoniae* the most active antibiotic was ertapenem (78%).

**Conclusions:** The antibiotic most active against all urinary Enterobacteriaceae isolates was ertapenem. Antibiotics commonly used to treat urinary tract infections, such as fluoroquinolones and trimethoprim sulfamethoxazol were less efficient, therefore their empirical use should be avoided. Nitrofurantoin, an antibiotic used to treat uncomplicated urinary tract infections caused by *E. coli*, was highly active.

**Keywords:** urinary isolates, fluoroquinolones, *E. coli*, *Klebsiella* spp.

## 282. A SYSTEMATIC LITERATURE REVIEW OF HEREDITARY ASPECTS OF OVARIAN CANCER

**Dumitrita Colac**

Scientific adviser: Rotaru Ludmila, MD, Molecular Biology and Human Genetics Department of *Nicolae Testemitanu* State University of Medicine and Pharmacy, Chisinau, Moldova

**Introduction:** Every year, worldwide, are registered 10 million new cases of cancer and 6.2 million deaths over the cancer. About 5 to 40 % of malignant tumors of all anatomical locations have a genetic etiology, and this percentage is growing due to increased general morbidity. So far, in the literature there are described over 200 hereditary cancer syndromes, for 35% of which are fully described the primary molecular defects or localisation of chromosomal mutation, and DNA diagnosis has become a routine method of investigation for genetic diagnosis. Ovarian cancer also refers to these pathologies.