

duration of up to one hour, the absolute risk of developing purulent septic complications is 173.9 cases per 1000 patients, if the surgery duration was from 1 to 2 hours, the risk of purulent septic complications is 341.1 cases per 1,000 operated patients, and surgery lasting more than 2 hours increased the risk of postoperative septic infection to 416.6 cases per 1000 of patients operated. According to the results obtained in the group of patients who underwent a single surgery the incidence of postsurgical septic complications are 192.1 cases per 1,000 patients, whereas in the group of patients who have undergone two surgeries this index is 700.0 cases 1000 patients operated, which is 3.62 times higher.

Conclusions: Length of hospital stay in patients with septic postoperative complications is about 2 times higher compared to the duration of hospital stay in patients without septic complications and is, average, 12.859 ± 1.934 days/bed . On average, each case of septic, nosocomial infection, made hospital stay of the patient longer up to 7.326 days, this led to significant additional expenditures.

Keywords: surgery, purulent septic infection, septic postoperative complication

44. THE INCIDENCE OF SEPTIC NOSOCOMIAL COMPLICATIONS IN SURGICAL WARDS

Prisacaru Ion

Academic adviser: **Ghidirim Gheorghe**, MD, Ph.D., Professor, Academician, IP "Nicolae Testemițanu" State Medical and Pharmaceutical University, Chișinău, Republic of Moldova

Introduction: Nosocomial infections continue to be a current issue for all medical institutions, especially in surgical profile.

Materials and Methods: We studied 462 observation forms, including 249 patients treated conservatively and 213 sheets of patients treated surgically. Were extensive and intensive use indices, absolute risk, risk attribution and the relative risk were used.

Results: The study found that of 462 patients hospitalized in the Department of aseptic surgery, 57 patients developed septic nosocomial infection, the incidence constituting 12.34%, or 123.4 cases per 1000 patients (‰). It was found that the nosocomial septic complication's actual incidence in patients treated conservatively is 3.6 % and, in the group of patients undergoing surgery, the real incidence by septic infections is 24.6%. Therefore, the absolute risk of making a septic complication, based on our study is: for patients treated conservatively - 36.14 % and among patients undergoing surgery - 246.2 %. Attributable risk of surgery for septic complications is 210.06 cases per 1,000 patients operated, and the relative risk in people exposed to surgery is 6.81. Therefore, patients undergoing surgery have about 7 times higher chance of septic complications compared with patients treated conservatively. Risk of septic complications turned out to be higher in men (44.0,%) compared with female patients (56.0 %), the incidence rate is reversed, and the incidence of septic complications is 142.85% for men and 108,1% for women.

Conclusions: The real incidence of septic nosocomial infections in aseptic surgery ward is 123.4% of hospitalized patients, including conservative treatment group, the risk of developing of septic complications is 36.0 % and in the group of patients, undergoing surgery is 246.2 %.

Key words: surgery, septic complication, nosocomial infection

45. IMMUNOHISTOCHEMISTRY PROFILE OF THE MUCINOUS LESIONS OF APPENDICEAL AND OVARIAN ORIGIN

Vozian Marin, Chemencedji Inga, Negru Anastasia

Academic advisers: **Mishin Igor**, M.D., Ph.D., senior scientific researcher – coordinator, Laboratory of Hepato-Pancreato-Biliary Surgery; **Rojnoveanu Gheorghe**, M.D., Ph.D., university professor, Chair of Surgery no.1 "Nicolae Anestiadi", State University of Medicine and Pharmacy "Nicolae Testemițanu", Chisinau, Republic of Moldova

Introduction: Pseudomyxoma peritonei (PMP) is a rare clinical entity, characterized by a significant amount of mucinous ascites associated by peritoneal mucinous implants. The most frequent localization of primary lesion are the appendix and ovaries (epithelial mucinous lesions with varying