

degree of teeth, the thickness of buccal and oral alveolar plates, the anatomy of dental roots, the cervical outline and the relative position of cementoenamel junction.

Conclusion: The anatomical form of interdental septa determines the thickness of the cribriform plates, fact that plays an important role in the pathogenesis of a specific pathology. The rarefaction of the radiologic design in the marginal region of the septa is an initial, very important sign of the periodontal disease. There are some features of the radiologic image of normal septa that represent their anatomical particularities, and they show no pathological changes.

Keywords: Normal interdental and interradicular septa, periodontal disease, cementoenamel junction

28. USING SONIC RETRO TIPS IN THE RETROGRADE ENDODONTIC TREATMENT

Scutelnic Vladimir, Bolun Radu

Academic adviser: **Fala Valeriu**, M.D., Associate Professor, State Medical and Pharmaceutical University "Nicolae Testemițanu", Chișinău, Republic of Moldova

Introduction: The retrograde endodontic treatment of periapical periodontal diseases in classical vision shows little chance of success. This is explained by the use of inappropriate tools, inadequate visibility, frequent postoperative complications and failures that resulted in the extraction of the tooth. This changed radically with the introduction of the microscope, of micro tools, sonic and ultrasonic tips and plug biocompatible materials.

Purpose and Objectives: To evaluate preliminary clinical results of the use of SONIC retro tips in retrograde endodontic treatment and estimate the surgical access to the root apex and capacity of retrograde cavity preparation.

Material and methods: The study included 10 patients who had periapical periodontal disease at one tooth, 10 teeth underwent apical resection. While the clinical and instrumental examination, orthopantomography and dentoparodontal radiography were made, computed tomography in case of necessity.

Results: Following clinical examination data, we have set limits of periapical pathology, the condition of neighboring teeth, the condition of mucosa. Access to the operative field was superior in 100% of patients. On postoperative contact radiographs, the length of retrograde obturation and the length of resected tip were counted. During the 6 months period after surgery, 1 from 10 patients was complaining of periodical pains and there was sensibility around the root tip projection on palpation. According to the clinical and radiographic criteria, in 9 patients have obtained good results and only one of 10 has failed.

Conclusion: The tips simplify the surgical access to the root apex and have a good capacity of preparation. Preliminary clinical results indicate excellent progress at 6 months postoperatively. With these tips intra and postoperative aspects in apical resection are considerably improved for doctors and patients.

Keywords: periapical periodontal disease, retrograde preparation

29. TMJ DYSFUNCTION IN PATIENTS WITH MIGRAINE.

Arnaut Diana

Academic adviser: **Lupan Ion**, M.D., Ph.D., Associate Professor, Faculty of Dentistry, Department of Oro-Maxillo-Facial Dentistry, University „Nicolae Testemițanu”, Chișinău, Republic of Moldova

Moldovanu Ion, M.D., Ph.D., Associate Professor, Department of Neurology, University „Nicolae Testemițanu”, Chișinău, Republic of Moldova

Introduction: Temporomandibular joint disorders, or TMD, is a constellation of the group of orofacial pain, that includes masticatory muscle, articular conditions or both. This reference summary