

O267

HIPERFUNCTIA PARATIROIDIANA ITINERAR CHIRURGICAL: 63 CAZURI OPERATE

Diaconescu MR *, Glod M *, Costea I *, Diaconescu Smaranda **

Universitatea de Medicina si Farmacie "Gr T Popa" Iasi, Clinica IV-a Chirurgie, Clinica V-a Pediatrie***

Introducere: Multiplele aspecte etiopatogenice si anatomo-clinice ale hiperparatiroidismului (HP): pri-mar (HPP), renal (HPR), familial, din NEM etc constituie o patologie care realizeaza o continua provocare. In afara fenotipului si simptomatologiei polimorfe, a noilor achizitii diagnostice si terapeutice, frapeaza contrastul dintre incidenta/prevalenta sindromului in crestere in tarile dezvoltate – mai ales pe seama observatiilor asimptomatice - si seriile limitate numeric sau cazurile izolate cu manifestari “istorice”, publicate in literatura natiunilor “in tranzit” sau subdezvoltate.

Material si metoda: Din 1986 in clinica noastra au fost operate 63 observatii de HP: 20 HPP – adenoame=17, carcinoame=2, paratiromatoza=1 si 43 HPR – HP secundar (HPS)=23 si tertiar=20. Am inregistrat 44 femei si 19 barbati (raport 2,3/1) cu limite de varsta 15 – 67 (medie 47) ani. Diagnosticul si indicatia chirurgicala au fost stabilite clinic prin prezenta suferintei renale – urolitiasis multipla sau recidivata in HPP – insuficienta renala cronica in regim de hemodializa in HPR, sindrom osos manifest – osteoporoză, dureri osoase, chisturi si fracturi, manifestari neuromusculare, psihonevrotice, digestive si cardiovasculare in ambele varietati. Datele de laborator au obiectivat valori anormale ale calcuiului seric total si ionizat, fosforului si fosfatazei alcaline si in special ale iPTH iar explorarile localizatoare au inclus ultrasonografia – mai putin concludenta in leziunile multiglandulare si mai recent scintigrafia cu 99mTc-tetraphosmin. Rezultate: Toate cazurile au fost operate practicandu-se exeresă simplă in 17 adenoame si “in bloc” cu lobul tiroidian ipsilateral in doua cancer (unul fiind o recidiva la 4 ani dupa indepartarea unui adenom), intr-un adenom chistic intratiroidian ca si in cazul de paratiromatoza (de asemenea recidiva dupa exercitarea extra muros a unui adenom). In observatiile de HPR au fost executate 24 parathyroidectomii subtotale (in 20 observatii reusindu-se exeresă standard a 3 si ½ glande, in rest indepartandu-se 3 sau doar 2 paratiroidi) si respectiv 19 parathyroidectomii totale (6 cu autotransplant glandular si 13 simple). Din considerente tactice sau pentru leziuni asociate explorarea/exeresă chirurgicală a fost extinsa la tiroida (29 cazuri) sau timus (20 cazuri). Examenul anatomopatologic a precizat diagnosticul final in toate observatiile. Rezultatele imediate si in timp au fost bune in special in HPP. Nu au fost hipocalcemii persistente chiar in cazul parathyroidectomii extinse dar am notat o paralizie recurrentiala, un hematoma la lojă si recidiva in doua cazuri de autotransplant antibrachial ca si cea a unui cancer la 4 ani dupa extirparea unui adenom (leziune nouă?). Concluzii: Parathyroidectomia – cu rafinamentele sale recente: minim invaziva, endoscopică sau asistata robotică – este singurul tratament eficace si definitiv in HPP si constituie o terapie simptomatica importanta, desi suboptimal, in cazurile de HPR (a caror tratament ideal este transplantul renal). Exercizele paratiroidiene trebuie practice de specialisti antrenati in aceasta chirurgie.

SURGICAL ITINERARY IN PARATHYROID HYPERFUNCTION: 63 CASES OPERATED ON

Background: Hyperparathyroidism (HP) is a constantly evolving entity with multiple clinical varieties i.e.: primary (HPP), renal (HPR), familial, in MEN etc, proteiform phenotype and symptomatology, continuous modernizing diagnosis and therapeutic methods and striking differences in epidemiology between developed nations and the 3rd world's or “in transition” countries. Material and methods: The study population comprised 63 patients with HP operated on from 1986 in our clinic. There were 20 cases with PHP (17 adenomas, 2 carcinomas and one parathyromatosis) and 43 cases with RHP (23 secondary and 20 tertiary). The series included 44 women and 19 men (ratio 2,3/1), aged 15-67 (range 47) years. As a rule the documentation of signs and symptoms as well as recording of the surgical indications were consistently thorough. The presence of multiple and recurrent urolithiasis in HPP and renal failure on hemodialysis in HPR as well as bone, muscular, neuropsychiatric, digestive and cardiovascular manifestation in both syndromes are constantly described. Laboratory data indicated abnormally levels of serum calcium, phosphorus, alkaline phosphatase but especially of the iPTH. Localisation procedures included ultrasonography less valuable for multiglandular lesions and recently 99mTc-tetraphosmin scan. Results: All the cases were operated on: 17 simple exeresis for adenomas and 4 “en bloc” resections together with the thyroid lobe for two carcinomas, intrathyroid cystic adenoma and parathyromatosis one case each. In HPR 20 patients underwent standard subtotal parathyroidectomy (3 and ½ glands) but in 3 cases only 3 or even 2 glands were found and 19 total parathyroidectomy respectively (6 with auto-transplantation). Thyroid (n=29) and thymus (20) resections were practiced for associated lesions or tactical reasons. Pathology established the final diagnosis. Immediate and late results were good especially in HPP. Persistent hypocalcemia was not encountered even in extended resections but we avow a cervical hematoma, a recurrent laryngeal nerve palsy, two antibrachial recurrences and a carcinoma developed four years after resection of an adenoma (new lesion?). Conclusions: Parathyroidectomy with its recent refinements in minimally invasive, endoscopic, video- and robotic techniques constitutes the gold standard therapy for HPP and still remains the only permanently effective method offering an improved quality of life in HPR. These operations must be done by high-specialised surgeons.

O268

NODULUL TIROIDIAN SOLITAR: PARADIGME DIAGNOSTICE SI TERAPEUTICE

Diaconescu MR *, Diaconescu Smaranda *

Universitatea de Medicina si Farmacie "Gr T Popa" Iasi, Clinica IV-a Chirurgie, Clinica V-a Pediatrie***

Controversele privind diagnosticul si atitudinea terapeutica in nodulul tiroidian solitar (NTS) sunt proportionale numeric cu incidenta/prevalenta la nivel mondial a acestei entitati constituind o continua provo-care pentru specialistii in domeniul. Discrepanța dintre frecventa importanta a NTS si numarul relativ redus al tumorilor maligne tiroidiene nu anuleaza teama de a ignora un cancer ceea ce conduce la o chirurgie maximalista sau chiar