

intravenously 0.3-0.4 mg / kg / hour ataractice (seduksen, valium) and 3-5 mg / kg / hour titrated morphine solution. Effectiveness of anesthetic effect was assessed by monitoring systolic, diastolic and mean frequency and heart rate, acid-base balance parameters and electrolyte monitoring of blood biochemical indices (blood glucose indices). According to our methodology in 2004-2010 were operated 103 patients, aged 54-84 years. Pirogov type mastectomy was performed in 48 patients, simple mastectomy in 21 patients, 34 patients extended sectoral resection. Intra- and postoperative period was devoid of features. 69 patients observed over 5 years, local recurrences were detected in 3 patients, distant metastases or not found. The results received show that the proposed techniques extend the indications for surgery in patients with BC in menopause.

E105

EFICACITATEA DIVERSELOR METODE DE INTERVENȚII CHIRURGICALE ÎN CADRUL TRATAMENTULUI COMBINAT ȘI COMPLEX

Godoroja Nadejda, Botnariuc Natalia, Cotruta A.

Institutul Oncologic, Republica Moldova

Prin analiza rezultatelor tratamentului a 496 bolnave de CGM st. I, 77 bolnave st. IIA (T2N0M0) și 295 bolnave st. IIB – IIIA (răspândire locoregională) s-a stabilit că în st. I creștere multicentrică nu a fost stabilită divergența semnificativă a frecvenței de progresare după operație organomenajantă (OOM) și mastectomie (ME) în cadrul tratamentului complex și combinat – corespunzător 11,4%; 3,6% și 17,9%; 5,9%. Diametrul tumorii mai mare de 3 cm în st. IIA mărește frecvența metastazării după OOM cu 5,2% în tratamentul combinat și 1,9% în tratamentul complex. Răspândirea locoregională (T1-3N1-2) a mărit frecvența progresării după tratament complex, varianta intervenției chirurgicale – ME până la 22,5% în T1-2N2M0 și 39,6% în T3N1-2M0. Depistarea precoce și implementarea CT și Rath neoadjuvante și adjuvante în cadrul tratamentului combinat sau complex mărește posibilitatea micșorării volumului intervenției chirurgicale.

EFFECTIVENESS OF VARIOUS METHODS OF SURGERY IN THE TREATMENT OF COMPLEX COMBINED

By analyzing the results of treatment of patients BC 496 st. I, 77 still. IIA (T2N0M0) and 295 ill st. IIB - IIIA (locoregional spread) has been established that in st. I multicenter growth has not been established significant divergence in the frequency of progression after surgery radical resection and mastectomy (ME) in the complex and combined treatment 11.4% accordingly, 3.6% and 17.9%, 5.9%. Tumor diameter greater than 3 cm in st. IIA increases the frequency Metastatic radical resection after combined treatment with 5.2% and 1.9% in the complex treatment. Locoregional spread (T1-3N1-2) progressing after treatment increased the frequency of complex variant surgery - ME to 22.5% in T1 and 39.6% in T3N1-2M0. Early detection and implementation of neoadjuvant and adjuvant CT and Rath in combination therapy can decrease the volume of complex surgery.

E106

SCHWANNOM GIGANT RETROPERITONEAL

Ghiur M., Ghiur Luminița, Serbanoiu D., Munteanu I.

Spit. CF 2

OBIECTIVUL STUDIULUI: Se prezinta cazul unei tumori retroperitoneale de schwannom gigant. Schwannoamele sunt tumori neurogene, non epiteliale ce se dezvoltă din teciile nervilor periferici (redenumite recent- PNST). Localizarea retroperitoneala a acestor tumori este foarte rar întâlnită în practica clinică (0,3-0,5%). Lucrarea prezintă cazul unui bolnav în vârstă de 52 de ani, de profesie inginer, fără alte antecedente patologice sau formațiuni tegumentare de neurofibromatoză care a fost spitalizat pentru dureri abdominale, mărirea de volum a abdomenului și creșterea cu 9 kg. Explorările imagistice și biologice care arată o tumoră retroperitoneală giganta, banuită benignă, nu au putut preciza originea tumorii. Se practica laparotomie mediană xifo-pubiană cu abord transperitoneal. Intraoperator se constată o tumoră retroperitoneală giganta de aspect lipomatos. S-a efectuat ablatia completă a tumorii fără sacrificii de organe. Dimensiunile foarte mari ale tumorii, greutatea de 8000 gr. și aspectul macroscopic sugerează benignitatea. Examenul histopatologic al tumorii este de schwannom benign, confirmat și de examenul imunohistochimic. Evoluție favorabilă în 14 zile postoperator. **CONCLUZII:** Spațiul retroperitoneal este foarte generos pentru dezvoltarea unor mase tumorale foarte mari. Incidența schwannoamelor este foarte mică. Prezentarea clinică a acestor tumori este nespecifică, cu dificultăți de diagnostic preoperator. Absența invaziei capsulare și a metastazelor, confirmă natura benignă a acestor tumori.

GIGANT RETROPERITONEAL SCHWANNOMA

THE OBJECTIVE OF THE STUDY: Case report of the tumor retroperitoneal of the giant schwannoma. The schwannomas are neurogenic tumors non epithelial which develop of the peripheral nerve sheaths (renamed recently- PNST). Retroperitoneal localization of this tumor is low in clinical practice (0,3-0,5%). Case report- an 52-year-old male patient, by engineer profession, without pathological records and without neurofibromatosis type tegumentary lesions, was hospitalized for abdominal pains, a change of the aspect of the abdomen and taken in weight 9 kilograms. Imaging and biologic tests which establish a giant retroperitoneal tumor, possibly benign, couldn't define the exact origin. In xifo-pubian laparotomy and transperitoneal abord we found a giant retroperitoneal tumor of fatty aspect. Tumor excision was performed without viscus sacrifice. The very big size of the

tumor, weight-8000gr., the macroscopic aspect suggests benign tumor. Histological examination of neurogenic tumor showed a benign schwannoma, confirmed also of the immunohistochemical test. Postoperative course was favorable in 14 days. CONCLUSIONS: The retroperitoneal space is gratly for the growth of the big tumors. The incidence of the schwannomas is low. The clinical presentation of these tumors is non specific, with difficulties for the correct diagnosis in preoperative period. The absence of the capsule invasion and of the metastasis confirme benign origin of these tumors.

E107

TUMORA STROMALA AGRESIVA CU LOCALIZARE GASTRICA- PREZENTARE DE CAZ

Anitei Maria-Gabriela, Scripcariu D. V., Scripcariu V.

Spital Sf Spiridon Iasi, Clinica III Chirurgie

Rezumat: Tumora stromala gastrointestinala (GIST) sunt definite printr-un ansamblu de argumente clinice, morfologice si imunohistologice. Prezentam cazul unei paciente in varsta de 39 ani cu GIST localizata la nivelul fornixului gastric cu metastaza hepatica, cu simptomatologie clinica nespecifica. Investigatiile paraclinice - ecografia abdominala si examenul computer tomografic (CT) - au evidentiat masele tumorale intraperitoneale fara a putea sugera diagnosticul. Examenul anatomo- patologic extemporaneu stabileste diagnosticul de tumora stromala gastrica si impune conduita chirurgicala corespunzatoare. Cuvinte cheie: GIST, metastaza hepatica

AGGRESSIVE GIST WITH GASTRIC LOCATION – CASE REPORT

Abstract: Gastrointestinal stromal tumors (GIST) are defined by a set of clinical, morphological and immunohistological elements. We present a patient, 39 years old, female, with GIST located in the upper part of the stomach with liver metastase, with nonspecific clinical symptoms. Paraclinical - abdominal ultrasound examination and computer tomography (CT) - have shown intraperitoneal tumor masses without suggested the diagnosis of GIST. Anatomic-pathological examination, made during the operation, diagnosed stromal tumors and gastric surgery requires proper conduct. Keywords: GIST, liver metastasis

E108

PARTICULARITATI IN TRATAMENTUL NEOPLASMULUI DE RECT JOS SITUAT

Anitei Maria-Gabriela, Giurgiu Iuliana, Grigoras M., Scripcariu V.

Abstract: Introducere Neoplasmul rectal beneficiază de tratament multimodal ce trebuie adaptat fiecărui caz în parte. Material și metode : Este prezentată experiența unei singure echipe pe un număr de 167 cazuri de neoplasm rectal tratate în perioada 2005-2011. Cazurile au fost supuse unui protocol de stadializare preterapeutică și s-au evidentiat 106 pacienți cu tumori situate în jurul distanței de 6 cm de marginea anală. Tratamentul neoadjuvant a fost reprezentat de radioterapie în regim de lungă durată pentru 89 de cazuri (T3, T4), iar chimioterapia a fost asociată în 56 cazuri. S-au practicat 19 rezecții anterioare de rect foarte joase cu excizie totală de mezorect, cu 11 anastomozes manuale și 8 anastomozes mecanice, 65 excizii abdomino-perineale de rect (18 asistate laparoscopice), 6 cazuri excizii locale transanale pentru tumori T0 și T1 și 16 operații Hartmann cu excizie totală de mezorect. Ileostomia de protecție s-a practicat în 14 cazuri de rezecții foarte joase de rect și a fost închisă la 6-8 săptămâni. Rezultate Complicațiile imediate postoperatorii au fost: fistule anastomotice 1, fistulă recto-vaginală 3, disfuncții urinare 4. Recidivele locale după rezecțiile anterioare foarte joase de rect au fost întâlnite la 3 cazuri iar după excizie abdomino-perineală la 2 cazuri. Concluzii : Rezultatele postoperatorii sunt dependente de o corectă stadializare. Pacienții cu tumori jos situate tratate prin excizie abdomino-perineală de rect au avut tumori mai mari și mai agresive decât pacienții tratați prin rezecții anterioare foarte joase de rect Cuvinte cheie : cancer rectal, rezecție anterioară de rect, excizie abdomino-perineală de rect

THE PARTICULAR PROBLEMS OF LOW RECTAL CANCER

Introduction: The gold standard in rectal cancer today is a multimodal approach, tailored for each patient. Material and methods: It is presented the one team experience with 167 cases of rectal cancer treated between 2005-2011. All patients have undergone a pretherapeutic staging protocol and there were 106 patients with tumours within 6 cm of anal verge. The neoadjuvant treatment was radiotherapy for 89 cases (T3, T4) and chemotherapy was associated in 56 cases. The procedures performed for low rectal cancer were 19 resections considered very low, with total mesorectal excision with 11 manual anastomosis and 8 mechanical anastomosis; 65 abdomino-perineal excision of the rectum (18 assisted laparoscopic), 6 transanal excision for T0 and T1 tumors and 16 extended Hartmann's procedure, with total mesorectal excision. Protective ileostomy has been practiced for 14 cases of very low resection of the rectum and was closed 6-8 weeks later. Results: Postoperative complications were: anastomotic leakage-1, recto-vaginal fistula-3, urinary dysfunction-4. Local recurrence after very low anterior resection of the rectum were diagnosed in 3 cases and after abdomino-perineal excision in 2 cases. Conclusion: It is extremely important to perform a preoperative staging of all the diagnosed cancer patient in order to get the best results. Patients with low rectal cancer treated by abdomino-perineal excision have worse tumours and higher involved margin rates compared with those treated by anterior resection. Keywords: low rectal cancer, anterior resection, abdomino-perineal excision