


**RESECTIONS IN ADVANCED PANCREATIC, COLON AND GASTRIC CANCER.**


Hotineanu A., Burgoci S.

State Medical University „N. Testemitanu”; Chisinau, Moldova. 2 Department of Surgery.


Introduction : The prognosis for life in patients with locally advanced malignant tumors is extremely poor in any primary location. The invasion of adjacent organs and trunk vessels over the decades has been considered unresectable for technical reasons and also for oncological radicalism. The aggressive surgical approach will extend the limits of the resection and in the association of perioperative chemotherapy can provide greater survival in this patient population.



Hemicolectomie dreaptă cu duodenpancreatectomie cefalică




Gastrectomie totală, ELG D2 Rezeție hepatică SII-III, Rezeție distală de pancreas cu prezervarea splinei



Pancreas, splina, stomac, colon - mobilizate



Cancerul de colon cu invazie în duoden și cefal pancreatic



Rectal cancer with invasion of the vagina, uterus and bladder




Corporeocaudal pancreatic cystadenocarcinoma invasion in the stomach, jejunum, colon, liver


Material and methods: We analyzed 54 patients with locally advanced malignant tumors with invasion in neighboring organs, which underwent multivisceral resections. From the total group 12 patients were diagnosed with gastric cancer, 24 patients with colorectal cancer, 18 patients with pancreatic cancer.

Results: R0 resection was achieved in 86% of cases. Postoperative morbidity 7.4%, morbidity 21.4% and are directly proportional to the number of anastomosis performed in stages of reconstruction.

Conclusions: The morbidity and mortality of multiorgan resections is significantly higher than standard resections, therefore requires careful selection of patients for aggressive surgical approach. Multivisceral resections have curative potential and are feasible in R0. Overall survival in locally advanced in IV stage for patients after multivisceral resection is much longer than patients after palliative treatment.



En Bloc Right Hemicolectomy with Pancreaticoduodenectomy for Locally Advanced Right-Sided Colon Cancer



Survival analysis after pancreatic resection for pancreatic cancer with multivisceral resection

Survival analysis after gastrectomy for stomach cancer with multivisceral resection