

OBSTRUCTIVE SARCOIDOSIS

Calaraş Diana, Munteanu Oxana

Academic adviser: Botnaru Victor, M.D., Ph.D., Professor, State Medical and Pharmaceutical University "Nicolae Testemiţanu", Chisinau, Republic of Moldova

Introduction: Sarcoidosis is a multisystemic granulomatous disease of unknown cause, that mainly affects the lungs. Being an interstitial lung disease, it is generally accepted that sarcoidosis demonstrates mostly a restrictive pattern in pulmonary function tests. Some authors mention that sarcoidosis could show an obstructive pattern also, but to a lesser extent.

Aim: to assess the frequency of the obstructive and restrictive patterns in patients with pulmonary sarcoidosis using pulmonary function tests.

Materials and methods: We performed an analysis of 95 patient's clinical records with biopsy proven or highly suggestive of sarcoidosis, cases registered in a third level medical institution, during year 2011. All the patients underwent chest X ray and pulmonary function tests.

Results: Our study group consisted of 74 (77,9%) females and 21 (22,1%) males, mean age $49,5 \pm 9,0$. In our group, 86 (90,5%) were non-smokers, ex-smokers - 4 (4,2%), and current smokers - 5 (5,3%). According to the radiological stages we had 25 (26,9%) subjects in stage I, 49 (52,7%) in stage II, 13 (14,0%) - stage III and stage IV - 6 (6,5%) patients. We found decreased FEV1 and decreased FEV1/FVC only in 8 (8,4%) cases, but we have also found other functional signs of obstruction, like decreased MMEF_{25-75%} in 51 (53,7%) cases, suggesting small airway obstruction; increased RV in 40 (42,1%) cases - corresponding to the air-trapping phenomenon, and increased TLC in 20 (21,3%) cases - suggesting hyperinflation.

In contrast, we found only 15 (15,7%) cases of concomitant decrease of FVC and normal or increased FEV1/FVC, suggestive of restrictive pattern, also decreased TLC in 8(8,5%) subjects, decreased RV in 4(4,3%) cases; concomitant decreased FVC, increased FEV1/FVC and decreased TLC - in 8 (8,5%) cases.

Conclusion: Our study showed that pulmonary sarcoidosis determines more obstructive defects than restrictive, depending on how we define obstruction.

Key words: sarcoidosis, obstruction, restriction, pulmonary function tests

CORRELATIONS COMORBIDITIES ON QUALITY OF LIFE IN PATIENTS WITH COPD

Brunchi Lucia, Butnaru Mihaela

Academic adviser: Alexandru Corlăţeanu, M.D., Ph.D., Senior lecturer, State Medical and Pharmaceutical University "Nicolae Testemiţanu", Chisinau, Republic of Moldova

Introduction: Chronic Obstructive Pulmonary Disease (COPD) is characterized by limited airflow that is incompletely reversible, can progress and may be associated with an abnormal inflammatory response of lungs to irritants. COPD is a major cause of morbidity in the elderly, affecting about 15% of the population aged over 65 years.

Material and Methods: In the questionnaires the following indices were studied: age, 6 minutes walk test (6MWD), index smoker - pack / year, Health - related quality of life (HRQL) instruments included: Saint George's Respiratory Questionnaire (SGRQ), Clinical COPD Questionnaire (CCQ), and comorbidities evaluated by Charlson and CDS index. The study was conducted on a group of 60 patients (30 men and 30 women) aged between 50-80 years and average of $64.45 (\pm 8.59)$.