

6. Turner D. et al. *C-reactive protein (CRP), erythrocyte sedimentation rate (ESR) or both? A systematic evaluation in pediatric ulcerative colitis*. In: *J. Crohn Colit*, 2011, vol. 5, p. 423-429.
7. Gomes P. et al. *Relationship between disease activity indices and colonoscopic findings in patients with colonic inflammatory bowel disease*. In: *Gut.*, 1986, vol. 27, p. 92-95.

Elina Berliba, d.ș.m, conf. univ.,
 Disciplina Gastroenterologie, Departamentul Medicină Internă,
 USMF Nicolae Testemițanu,
 Tel.: 079689636
 E-mail: alina_berliba@yahoo.com

ULCERATIVE COLITIS AND CYTOMEGALOVIRUS INFECTION

Alina JUCOV, Svetlana ȚURCAN,
 State University of Medicine and Pharmacy
 Nicolae Testemițanu, Chișinău, Moldova

Rezumat

Colita ulceroasă și infecția cu citomegalovirus

Boala inflamatorie intestinală (BII) este numele generic a două entități: boala Crohn și colita ulceroasă care reprezintă o inflamație cronică nespecifică a tractului gastrointestinal. Infecția cu citomegalovirus (CMV) este o infecție virală comună la om. Ea determină de multe ori infecție primară și ulterior persistă pe tot parcursul vieții într-un stadiu latent. În diverse situații de imunosupresie, virusul se poate reactiva și cauza boli, afectând multiple organe, inclusiv tractul gastrointestinal. Etiologia BII nu este clară și infecția CMV este deseori asociată cu BII. În acest articol este analizată legătura patogenetică dintre BII și infecția CMV.

Cuvinte-cheie: *boală inflamatorie intestinală, colită ulceroasă, boala Crohn, citomegalovirus*

Резюме

Язвенный колит и цитомегаловирусная инфекция

Воспалительные заболевания кишечника (ВЗК) – это общее название хронических неспецифических поражений кишечника, которые включают язвенный колит и болезнь Крона. Цитомегаловирусная (ЦМВ) инфекция, с другой стороны, представляет собой распространенное вирусное заболевание, которое зачастую обуславливает первичную инфекцию и затем персистирует в организме в неактивной форме. При развитии иммуносупрессии, вирусная инфекция может активироваться и вызывать различные патологии, включая поражение желудочно-кишечного тракта. Этиология ВЗК неизвестна, однако отмечена частая

ассоциация ВЗК с ЦМВ инфекцией. В данном обзоре литературы обсуждаются возможные патогенетические связи между ВЗК и ЦМВ.

Ключевые слова: *воспалительные заболевания кишечника, язвенный колит, болезнь Крона, цитомегаловирусная инфекция*

Introduction

Inflammatory bowel disease (IBD) represents chronic non-specific inflammation of gastrointestinal tract. The etiology of IBD is not clear and Cytomegalovirus infection is often associated with IBD. Early studies indicated that CMV infection can lead to subsequent development of IBD [1]. This may be possible in some susceptible patients but in the most recent reports CMV colitis occurred primarily in patients with pre-existing IBD [2, 3]. The pathogenetic link between IBD and CMV infection was supposed and began to be studied in the last decades.

The aim of the study was to evaluate critically literature data on the relationship between CMV and IBD.

Methods

Internet search was conducted in Medline (from 1966 to 2013) and PubMed (from 1980 to 2013) database using words "cytomegalovirus", "inflammatory bowel disease", "ulcerative colitis", "Crohn's disease".

Results and discussion

419 articles were found. The prevalence of CMV infection is high, ranging from 30–100%, depending on age and race. Cytomegalovirus is a member of the Herpesviridae family which contains a double-stranded DNA. Most CMV infections are acquired either in the perinatal period and infancy or in adulthood through sexual contact [4]. It often causes primary infection in humans, and later persists lifelong in a latent stage. Primary infection in the immunocompetent hosts is usually asymptomatic or causes syndrome with fever, myalgia and pharyngitis. Gastrointestinal affectation due to primary infection has been described but is rare, and manifests with bloody diarrhea, tenesmus, abdominal pain, fever, anorexia, malaise and weight loss [3]. CMV can cause esophagitis, gastritis, ulcers, terminal ileitis and colitis.

In latent infection principal reservoirs of CMV are fibroblasts, myeloid cells and endothelial cells [5]. Peripheral blood monocytes constitute a major site of viral latency and may be triggered by secreted proinflammatory cytokines and chemokines. They can differentiate into tissue macrophages, leading to CMV reactivation and probably to CMV disease. Endothelial cells are also a common target for CMV

in vivo, regardless of the organ involved. The vascular endothelium represents the interface between circulating immune cells and the lamina propria of the gut and this can partially explain the role of CMV in cases of intestinal inflammation.

The association between CMV and IBD was described long ago. The first case report dates from 1961, when Powell and coauthors described a patient with UC and "cytomegalic inclusion" disease. Since then, questions about the role of CMV in these patients remains open [7]. Does CMV reactivation exacerbate the disease in patients with pre-existing IBD or is reactivation a consequence of IBD activity and its treatment with CMV acting as an innocent bystander [3, 8]? Interpretation of existing results is limited because most studies are small and retrospective, with different diagnostic methods of CMV detection and even different classifications of the severity of concomitant IBD.

Patients with IBD are immunosuppressed due to chronic inflammation, medications and poor nutrition [2, 9]. CMV reactivation may be triggered by TNF- α , catecholamines and proinflammatory prostaglandins [10]. During active IBD, local expression of a wide variety of cytokines including TNF- α , IFN- γ , and IL-2 is induced, with activation of transcription factors (NF-Kb) and production of chemokines and adhesion molecules that recruit circulating monocytes and dendritic cells in the area of inflammation. There, these cells differentiate into permissive cells supporting active replication of the virus. Endothelial cells can also serve as permissive cells as they have been shown to stimulate T-cells to produce IL-2 and to proliferate. Activated T-cells consequently produce TNF- α and IFN- γ and perpetuate the inflammation process causing more injury to the gut. These indicate that CMV has tropism for sites of inflammation and confirm the results of clinical studies that have shown presence of CMV in the affected region of the gut in the IBD patients. They also indicate that CMV replication is the result of CMV reactivation rather than primary infection.

Experimental studies have identified three factors that influence the reactivation of CMV infection in active colitis:

- increased cell proliferation in inflamed tissue with ulcers that attract CMV;
- inherent impaired natural killer cell activity presented by patients with IBD;
- use of immunosuppressive drug [6].

The third factor is controversial, since use of steroids may be either a risk factor or a surrogate marker of severity disease [2, 9]. In vitro data suggest that steroids and cyclosporine could support the replication of CMV [11]. Recent studies suggested

CMV can appear only in inflamed tissue and is not found in healthy tissue.

Hommel and coauthors proposed the following sequence to explain the pathophysiology of CMV disease in patients with active UC:

- initiation phase, mucosal inflammatory response induces expression of cytokines and chemokines which activates latently infected cells and the migration of monocytes and dendritic cells into the inflamed mucosa;
- reactivation phase, in which infected monocytes differentiate into tissue macrophages and dendritic cells;
- consolidation phase, during which the virus causes an active replication predominantly in endothelial cells that likely exacerbates inflammation [10].

Theoretically, all latently infected IBD patients receiving immunosuppressive treatment frequently produce infectious virus and don't allow antiviral immune responses to develop. In two recent clinical studies, corticosteroids did not seem to be a major factor in the development of CMV infection and disease in IBD patients [12, 13].

There have also been reports of colitis patients with evidence of active CMV infection who improved with steroids and did not require antiviral treatment [14], as well as patients with active colonic CMV infection without active colitis [15]. In these cases CMV seems to behave like an innocent bystander. As it has been mentioned before, CMV has the propensity to infect rapidly growing tissue, especially endothelial cells in granulation tissue. Some studies suggested that CMV represents an opportunistic infection in severely inflamed mucosa rather than a primary pathogen [16].

The most widely held theory is that CMV infects areas of active IBD and causes further tissue injury aggravating the severity of the underlying IBD. In the majority of case-reports patients with severe attacks of IBD and CMV infection had significant morbidity (toxic megacolon 15%, colectomy up to 62%) and mortality (up to 44%). Antiviral treatment prevented colectomy in some but not all of the patients. In more recent series the mortality rate of CMV colitis in UC were 30% and the rate of surgery 40% [9]. CMV disease seems to be less frequent in patients with Crohn's disease compared to patients with ulcerative colitis.

Conclusion

The role of CMV infection in patients with IBD has not yet been clearly defined. In the majority of published studies CMV is considered to act like a true pathogen, complicating the course of IBD, causing

the resistance while in the others, CMV does not seem to alter the natural course of the underlying IBD. Further prospective studies are needed to clarify the role of CMV infection

References

1. Orvar K., Murray J., Carmen G. et al. *Cytomegalovirus infection associated with onset of inflammatory bowel disease*. In: Dig. Dis. Sci., 1993, nr. 38, p. 2307–2310.
2. Papadakis K.A., Tung J.K., Binder S.W. et al. *Outcome of cytomegalovirus infections in patients with inflammatory bowel disease*. In: Am. J. Gastroenterol., 2001, nr. 96, p. 2137–2142.
3. Domènech E., Vega R., Ojanguren I. et al. *Cytomegalovirus infection in ulcerative colitis: a prospective, comparative study on prevalence and diagnostic strategy*. In: Inflamm. Bowel Dis., 2008, nr. 14, p. 1373–1379.
4. De la Hoz R. E., Stephens G., Sherlock C. *Diagnosis and treatment approaches of CMV infections in adult patients*. In: J. Clin. Virol., 2002, nr. 25 (suppl. 2), p. S1–12.
5. Plachter B., Sinzger C., Jahn G. *Cell types involved in replication and distribution of human cytomegalovirus*. In: Adv. Virus Res., 1996, nr. 46, p. 195–261.
6. Ayre K., Warren B.F., Jeffery K., Travis S.P. *The role of CMV in steroid-resistant ulcerative colitis: A systematic review*. In: J. Crohn's Colitis, 2009, nr. 3, p. 141–148.
7. Powell R.D., Warner N.E., Levine R.S., Kirsner J.B. *Cytomegalic inclusion disease and ulcerative colitis: report of a case in a young adult*. In: Am. J. Med., 1961, nr. 30, p. 334–340.
8. Lwawlor G., Moss A.C. *Cytomegalovirus in inflammatory bowel disease: pathogen or innocent bystander?* In: Inflamm. Bowel Dis., 2010, nr. 16, p. 1620–1627.
9. Kishore J., Ghoshal U., Ghoshal U.C. et al. *Infection with cytomegalovirus in patients with inflammatory bowel disease: Prevalence, clinical significance and outcome*. In: J. Med. Microbiol., 2004, nr. 53, p. 1155–1160.
10. Hommes D., Sterringa G., van Deventer S. et al. *The pathogenicity of cytomegalovirus in inflammatory bowel disease*. In: Inflamm. Bowel Dis., 2004, nr. 10, p. 245–250.
11. Widmann T., Sester U., Gärtner B.C. et al. *Levels of CMV specific CD4 T cells are dynamic and correlate with CMV viremia after allogeneic stem cell transplantation*. In: PLoS One, 2008, nr. 3, p. e3634.
12. Pfau P., Kochman M.L., Furth E. et al. *Cytomegalovirus colitis complicating ulcerative colitis in the steroid-naive patient*. In: Am. J. Gastroenterol., 2001, nr. 96, p. 895–899.
13. Martin S.I., Sepehr A., Fishman A.J. *Primary infection with cytomegalovirus in ulcerative colitis*. In: Dig. Dis. Sci., 2006, nr. 51, p. 2184–2187.
14. Criscuoli V., Casa A., Orlando A. et al. *Severe acute colitis associated with CMV: A prevalence study*. In: Dig. Liver Dis., 2004, nr. 36, p. 818–820.
15. Maconi G., Colombo E., Zerbi P. et al. *Prevalence, detection rate and outcome of cytomegalovirus infection in ulcerative colitis patients requiring colonic resection*. In: Dig. Liver Dis., 2005, nr. 37, p. 418–423.
16. Eyre-Brook I.A., Dundas S. *Incidence and clinical significance of colonic cytomegalovirus infection in idiopathic inflammatory bowel disease requiring colectomy*. In: Gut, 1986, nr. 27, p. 1419–1425.

PARTICULARITĂȚI CLINICE ȘI PARACLINICE ALE URTICARIEI CRONICE RECIDIVANTE, ASOCIATE CU PATOLOGIA GASTRODUODENALĂ

Vera ONU¹, Vlada-Tatiana DUMBRAVA¹,
Liudmila RUSSU¹, Marina CASAPCIUC², Iraida
SEREBREANSCHI², Natalia PISCIALNICOVA²,

¹Departamentul Medicina Internă, Clinica medicală nr. 4,
USMF Nicolae Testemițanu, ²Spitalul Clinic Republican

Summary

Clinical and paraclinical particularities in recurrent chronic urticaria associated with gastroduodenal pathology

The aim of the study was to examine the clinical and paraclinical particularities in patients with chronic recurrent urticaria (CRU) depending on the functional state of the gastrointestinal segment of the digestive tract.

It was performed a retrospective study based on clinical observation sheets in 60 patients diagnosed with chronic recurrent urticaria hospitalized in Republican Clinical Hospital in Allergology Department during 2010-2012. The results of instrumental and laboratory investigations to determine the clinical and paraclinical manifestations of CRU associated with gastroduodenal pathology were analyzed.

Study determined that CRU allergic and that associated with gastroduodenal pathology shown by similar cutaneous events (rash, pruritus), but with some differences between manifestations of urticarial phenomena, expansion area and evolution of damage gastroduodenal disease. In patients with CRU and urticaria associated with gastroduodenal pathology, urticarial process has pseudo-allergic character and as provocative factors serve the background disease, errors in diet, use of drugs, alcohol, stress. In patients with chronic recurrent urticaria in association with gastroduodenal pathology significantly is detected more frequently H. pylori infection compared to patients with chronic allergic urticaria. In patients with CRU associated with gastroduodenal pathology gastroduodenal disorders are directly correlated with the severity of urticaria evolution.

Keywords: *chronic recurrent urticaria, gastroduodenal pathology*

Резюме

Клинические и параклинические особенности хронической рецидивирующей крапивницей, ассоциированной с гастродуоденальной патологией

Цель исследования – изучение клинических и параклинических особенностей у больных с хронической рецидивирующей крапивницей (ХПК) в зависимости от функционального состояния желудочно-кишечного сегмента пищеварительного тракта.

Было проведено ретроспективное исследование на основании историй болезни 60 пациентов с диагнозом ХПК, госпитализированных в отделение Аллергологии